




P. Michael Link
Jasmin Kominek
Jürgen Scheffran

*Impacts of sea level rise
on the coastal zones of Egypt*

University of Hamburg
Research Group Climate Change and Security

Working Paper
CLISEC-25



Impacts of sea level rise on the coastal zones of Egypt

Working Paper CLISEC-25

P. Michael Link ^{a,b,c}, Jasmin Kominek ^{a,d} & Jürgen Scheffran ^{a,c}

^a *Research Group Climate Change and Security, KlimaCampus, University of Hamburg, Hamburg, Germany*

^b *Research Unit Sustainability and Global Change, Center for Marine and Atmospheric Sciences, University of Hamburg, Hamburg, Germany*

^c *Institute of Geography, University of Hamburg, Hamburg, Germany*

^d *Institute of Sociology, University of Hamburg, Hamburg, Germany*

Correspondence to P. Michael Link:

Tel. +49 (40) 42838-7719

Fax +49 (40) 42838-9211

eMail: michael.link@zmaw.de

Abstract

The coastal zones of Egypt are expected to be particularly vulnerable to the rise of the sea level in the Eastern Mediterranean. Depending on the scenario of sea level rise, a considerable share of the precious agricultural land in Egypt is threatened, likely to cause not only economic losses but also risks to the livelihoods of millions of inhabitants of the coastal zones and subsequently even possible dangers to the stability of the entire society. In this paper we look at the risks imposed by sea level rise on the Egyptian coastal zones and assess the adaptive capacity of the affected society to address these issues adequately. This analysis is conducted in the context of an analytical framework that links climate change and its implications to societal stability and conflict via interactions with aspects of resource availability and individual wellbeing.

1 Introduction: Sea level rise and its extent in the 21st century

In addition to rising temperatures, changing precipitation patterns, and increasing frequencies of severe weather events, the rise of the global mean sea level is considered to be one of the most dangerous consequences of climate change. This aspect of climate change is of particular interest as already more than one tenth of the world's population lives within 10 m elevation of the current sea level (McGranahan et al., 2007), which is potentially at risk of needing relocation as the sea level continues to rise. Furthermore, the tremendous rate at which coastal zones are developed also increases the vulnerability of these areas. To preserve the livelihoods of the inhabitants of the coastal areas and to protect the assets in these regions, it is necessary to have a profound understanding of the processes leading to sea level rise. This is important to effectively plan and manage adaptation options to deal with an increasing sea level.

The extent of sea level rise is influenced considerably by several factors. First of all, there is steric sea level rise, i.e. the thermal expansion of the oceans as a consequence of global warming. Another component is eustatic sea level rise due to the melting of glaciers and ice sheets in Greenland and the Antarctic, which store vast amounts of fresh water (Jacob et al., 2012). Furthermore, there are other terrestrial sources of water such as high mountain glaciers and rivers. All these contributions taken together affect the relative sea level rise. In

order to obtain estimates of the absolute sea level rise, the dynamics of ocean currents as well as the tectonic movement of the land arising from e.g. subsidence or uplift also have to be taken into consideration (Nicholls et al., 2011; Rahmstorf, 2012). The latter contribution is referred to as isostatic sea level rise.

Simulations based on the Special Report on Emission Scenarios (SRES)-of the Intergovernmental Panel on Climate Change (IPCC) have led to estimates of an increase in relative global mean sea level (Fig. 1) between 18 and 59 cm by the end of the 21st century (Intergovernmental Panel on Climate Change, 2007). Considerable uncertainties arise because the extent of the contribution of the melting of the Antarctic and Greenland ice sheets is unclear. This factor may increase substantially in importance beyond a certain tipping point in global temperature increase (Church et al., 2010). Recent assessments of the development of the sea level indicate that the actual increase in sea level is likely to be much closer to the upper bound of the projections than to the lower bound (Rahmstorf et al., 2007) (Schaeffer et al., 2012). Consequently, sea level rise can be expected to become an increasingly urgent issue among the consequences of climate change.




Figure 1: Projections of global mean sea level change. The grey areas indicate the uncertainty in the long term estimates of the rate of sea level change. The line is a reconstruction based on measurements using tide gauges (Intergovernmental Panel on Climate Change, 2007).

However, there is substantial regional differentiation when it comes to the amount of sea level rise. It is the global average change of the sea level with regional variations due to the ocean dynamics (Stammer, 2008) in combination with local land motion that actually determines whether a particular coastal zone is at risk. One of the regions that can be considered to be exceptionally vulnerable to the impacts of sea level rise is the Egyptian coast along the Eastern Mediterranean. This is because of an unfavorable combination of relative sea level rise and land movement in this region.

In this paper, we have a closer look at the impacts of sea level rise on this particular coastal region, which encompass ecological changes and land losses as well as the resulting economic and social implications. All of these interact in a complex way and consequently have to be considered jointly (c.f. figure in Scheffran et al., 2012a). In this context, we will also address the strategies and measures that already have been taken or need to be taken to counter the adverse impacts of sea level rise on Egypt.

2 The coastal areas of Egypt

The coasts of Egypt extend over more than 3500 km along the Eastern Mediterranean and the Red Sea. The Mediterranean coast of Egypt can be divided into four main subregions (Frihy and El-Sayed, 2012): The northwest coastal sector extends from Sallum to Alexandria. It can be characterized as being particularly suitable for recreational activities and tourism. The Alexandria coastal sector extends further eastward from Hammam to Abu Qir. Then, the Nile Delta coastal sector extends all the way to Port Said in the East. The easternmost sector of the Mediterranean coast of Egypt is the North Sinai coastal sector, which extends from Port Said to Rafah. This sector consists of large dunes that can reach considerable heights, thus naturally protecting the coastal area.

More than 32 million people live along the coastlines of the country, mostly in the Nile River Delta between Alexandria and Port Said. The shoreline of the Nile Delta itself is 275 km long and consists of several promontories that are separated by saddles and bays (Frihy, 2003). There are three large lagoons that are separated from the sea. Initially, seven branches of the Nile River formed the delta. By now, these have silted up and Nile water passes through two remaining branches, the Rosetta branch in the West and the Damietta branch in the East.

The Egyptian coast along the Mediterranean is intensively used by the Egyptian people. There is considerable tourism infrastructure as the coastal zones of Egypt are the most popular tourism destinations in the country (El Raey, 2011). Revenues from tourism amount to more than 11% of the Egyptian GDP. Furthermore, there is substantial agricultural use of the flat land in the Nile Delta for food production and of the lakes and lagoons for fisheries, the latter providing more than 60% of the total catch of Egyptian fishermen. Even though the area of the Nile Delta amounts to only four per cent of the entire land area of Egypt, this region contributes more than one third of Egypt's agricultural production and almost two thirds of the country's fisheries' output (Frihy, 2003). To add to the value of this coastal land, the Nile Delta is one of the few regions of Egypt in which large scale agricultural production is possible at all.

In the recent past, the Nile Delta has experienced considerable changes in environmental conditions. Some of them have been of natural, some of anthropogenic origin (Becker and Sultan, 2009; Syvitski et al., 2009). For instance, there is erosion along the delta coast (El Banna and Frihy, 2009; Frihy et al., 2008) and the drying up of wetlands (El-Asmar and Hereher, 2011), which occurs during expansions of agricultural production areas. Furthermore, there are considerable human-induced changes to the coastal morphology in the region of the Nile Delta. Land reclamation projects have added several hundreds of square kilometers of agricultural land (El Banna and Frihy, 2009), and previously unproductive land has been converted to fish farms, considerably increasing the economic output in this part of the country. These changes increase the vulnerability of the coastal zone with regard to sea level rise as natural protection is removed and the likelihood of subsidence of the land is increased. Therefore, protective measures become more and more important in order to avoid loss of land due to inundation even at moderate rates of sea level rise.

In this context, it has to be noted that the regional implications of sea level rise for the Egyptian coast vary depending on the geographic location. Assessments of historic records indicate that there is a continuous relative increase in sea level irrespective of changing climatic conditions that fluctuates between 1.8 and 4.9 mm/a (Frihy et al., 2010). The highest rates generally occur near the Rosetta promontory while there is less sea level rise near the harbor of Alexandria. These differences can be attributed to the local geological settings. Recent measurements at the tide gauge station in Alexandria using global positioning systems (GPS) yield a relative sea level rise of 1.7 mm/a (Shaker et al., 2011). Taking into account tectonic settling of the area at the rate of -0.47 mm/a, the measurement-based estimate of absolute sea level rise at the northwest Mediterranean coast of Egypt is almost 2.2 mm/a. This corresponds well with tidal gauge measurements at different sites that indicate an increase in water level in the past decades between 1.0 mm/a in Burullus and 2.3 mm/a in Port Said (Frihy, 2003).

Nonetheless, these current rates of eustatic sea level rise that have been measured do not yet pose a threat to the Nile Delta (Hereher, 2010). There are coastal sand dunes, ridges and elevated strips along the coast that serve as initial barriers. Furthermore, there are already active coastal protection measures such as seawalls and the elevated design of the international coastal highway that protect the delta against the sea. On the other hand, there are processes that lead to the submergence of parts of the delta. These include the removal of coastal dunes and subsidence from water pumping from fossil aquifers, which can reach up to 5 mm/a (Stanley, 1998) and occur because of tectonics and sediment compaction.

Such land subsidence, which shows the highest rates in the area of Alexandria, is an important issue. Subsidence appears to occur mainly during and after major earthquakes and therefore mainly in non-continuous fashion. But it has been accounted for in the past. Major tectonic activity has already been confirmed in the fourth century A.D. (Frihy, 2003). In combination with the low elevations and flat character of large parts of the coastal areas along the Mediterranean coast of Egypt, subsidence can pose considerable problems should the sea level continue to rise. Assessments with digital elevation models indicate that almost 20% of the Nile River Delta already lies below the mean sea level (Hereher, 2010) and that another 25% lies below an elevation of 2 m. All of these areas are prone to flooding if the sea level rose as projected.

Because of the diverse use of the coastal zones of Egypt, it is already necessary to protect the coast against the current moderate increase in sea level. However, there would be widespread consequences if the sea level rose at a considerably higher rate in the Eastern Mediterranean as this would pose an additional burden on the protective measures that may overstretch the adaptive capacity of the Egyptian society. Such impacts are not only confined to land losses through inundation and erosion but also affect the population of this area in many direct and indirect ways.

3 Ecological and economic consequences of sea level rise in Egypt

When looking at the implications of sea level rise for the Egyptian coastal zones, it is important to note that it is a combination of several factors that contribute to the vulnerability of the regions. Of course, one key factor is the extent of increase in the sea level due to changing climatic conditions (Frihy, 2003). But additionally, the tectonic setting of the individual parts of the coastline also has to be addressed: does the land subside or does it currently emerge? And how do human activities influence the local development of the coastal zones as large scale removal of water from ground water aquifers contributes significantly to land subsidence? Furthermore, short term meteorological events such as storm surges also have an impact on the coastal zones that should not be disregarded.

Assessments of the vulnerability of the Mediterranean coastline of Egypt that take into account a combination of the factors mentioned above indicate that approximately 30% can be considered to be vulnerable to sea level rise (Frihy, 2003). These areas mainly consist of narrow natural sand barriers, the low and flat coastal plain and some low-lying beaches consisting of very fine sand. In contrast, about 55% of the coast is naturally protected and thus fairly safe from sea level rise. The remaining part of the coast (15%) is artificially protected and continuously needs to be protected this way to maintain the integrity of the Egyptian Mediterranean coast.

Table 1: Vulnerabilities of the Egyptian Mediterranean coasts with regard to sea level rise (adapted from Frihy and El-Sayed, 2012)

sector	vulnerability
Northwest coast: coastal plain	vulnerable only in extreme events or safe
Northwest coast: low-lying beaches	highly vulnerable
Alexandria: coastal plain	vulnerable only in extreme events or safe
Alexandria: low land	vulnerable only in extreme events
Alexandria: low-lying beaches	highly vulnerable
Nile Delta: around Rosetta extension	moderately or hardly vulnerable
Nile Delta: west of Damietta port	moderately or hardly vulnerable
Nile Delta: coastal lagoon barrier	highly or moderately vulnerable
Nile Delta: sand dunes	safe
Nile Delta: accretional beaches	safe
Nile Delta: low-lying beaches	highly vulnerable
North Sinai: coastal plain	vulnerable only in extreme events or safe
North Sinai: Bardawil Lagoon	hardly vulnerable
North Sinai: flash flood drains	vulnerable only in extreme events
North Sinai: low-lying beaches	highly vulnerable

A vulnerability ranking of the different geographic sectors of the Egyptian Mediterranean coast indicates that in the northwestern coastal sector, the low-lying beaches are particularly vulnerable to sea level rise (Tab. 1) (Frihy and El-Sayed, 2012). The same holds for the region of Alexandria. In the Nile Delta region, also the coastal lagoon barriers are extremely vulnerable. All these are subject to subsidence or high rates of coastal erosion. The moderately vulnerable zones are naturally protected, whereas the artificially protected areas can be considered to now have a low vulnerability to the current rate of sea level rise.

When the sea level rises, increased storm surges become a more prominent threat to the coastal areas. An assessment of the impacts of these climate change impacts indicates that the area of the coastal zones in Egypt affected by storm surges would almost double as a consequence of higher storm surge intensity (Dasgupta et al., 2009b). More than 2.6 million people would then be additionally affected by increased inundation, with an emphasis on the region around Port Said, the urban area that is considered to be the most adversely affected city in the Mediterranean North African region with respect to the consequences of sea level rise (Elsharkawy et al., 2009).

In addition to the impacts of more extensive extreme events, inundation of the flat and low laying coastal land is the prime danger arising from sea level rise. Simulation results indicate that the Nile River Delta is one of the regions that would be most affected worldwide by a rise in sea level of one meter (Dasgupta et al., 2011). Vast areas of the directly exposed parts of the coastal zone would be flooded. If the sea level only rose by 0.5 m, more than 1800 km² of agricultural land would be lost (Fig. 2), affecting approximately 3.8 million people (FitzGerald et al., 2008). If the sea level rose by a whole meter, the number of people that would need to leave their land would increase to more than 6 million (Dasgupta et al., 2009a), as approximately 4500 km² of land become uninhabitable. But because of the heterogeneous terrain in the Nile Delta, not the entire plain is affected by sea level rise in the same way.

However, continued sea level rise also increases the risk of coastal erosion and saltwater intrusion, which are the other key impacts in addition to direct land losses (Frihy et al., 2010).




Figure 2: The land areas in the Nile Delta that would be flooded if the sea level rose of 0.5 m and if the sea level rose 1.0 m (FitzGerald et al., 2008): The Mediterranean is shown in dark grey, the areas affected by an increase in sea level of 0.5 m are shown in medium grey, and the areas affected by a sea level rise of 1.0 m are shown in light grey.

Salt water intrusion into ground water aquifers is a severe problem particularly in those areas of the coastal zones, which have an elevation that is only slightly above the current sea level. Model simulations suggest that an increase in sea level of 0.5 m leads to an intrusion of salt water into the Nile Delta aquifers, which could reach up to nine kilometers inland (Sherif and Singh, 1999). Subsequent studies have confirmed this order of magnitude (Werner and Simmons, 2009). Degeneration of fresh water sources over such a large area is likely to have a substantial negative impact on agricultural production in this area. Additionally, this implies a considerable challenge to provide the large population of Egyptians that live in the coastal zones with adequate amounts of clean potable water. Should it become impossible to sustain the population currently living near the Egyptian coasts, large numbers of coastal inhabitants would need to migrate into other parts of the country.

However, the most direct economic impacts of sea level rise on Egypt arise from losses of agricultural land due to inundation. Depending on the extent of losses of agricultural land in the Nile Delta, the economic consequences can vary considerably. If only the northernmost coastal areas in the Nile Delta are affected, the economic impact may remain limited (Frihy and El-Sayed, 2012). But for a more pronounced sea level rise the effects may become particularly devastating to the Egyptian economy since the Nile Delta is one of the few parts of the country that can be used for large scale agricultural production. Although coastal lagoons are expected to be able to adjust to salinity changes in environmental boundary conditions (Frihy, 2003), declines in fish catches in the lakes and lagoons in northern Egypt are likely, which in turn leads to the abandonment of the fisheries by many fishermen and the subsequent decline in Egyptian GDP (Elsharkawy et al., 2009).

Assessments of the economic consequences of sea level rise highlight the necessity of artificial coastal protection in Egypt on a large scale because even the economic losses incurred by an increase in sea level of only 0.5 m without any adaptation measures would be considerable. More than 1.5 million people only in the vicinity of the city of Alexandria would have to be relocated (El Raey, 1997). The damage to land and infrastructure, e.g. due to the abandoning of the harbor of Alexandria would amount to more than 30 billion US-Dollars.

But even with considerable artificial protection, the Egyptian economy would be seriously affected by an extensive rise in sea level (Dasgupta et al., 2011). If the sea level rose by one meter, the gross domestic product would decline by almost 6.5% with the agricultural sector being hit disproportionately strong with a decline by 12.5%. Such a development is a direct threat to food security for the rapidly growing population, as the needs for fundamental food crops could be fulfilled to a far lesser degree than at present.

However, agriculture and fisheries are not the only economic sectors that would be adversely affected by sea level rise. Particularly the western part of the Mediterranean coast of Egypt has experienced a considerable development of tourism infrastructure and recreational activities. Assessments indicate that sea level rise is likely to cause local erosion and inundation of sandy beaches, which would have negative implications for the affected resorts and tourist destinations (Frihy and El-Sayed, 2012). These impacts can be expected to be very serious although no assessments quantifying these consequences have been carried out yet. Similarly, there are possible impacts of sea level rise on oil and gas exploration in the Nile Delta, on road infrastructure (e.g. the coastal road from Rafah to Matruh), and on coastal ecosystems and wetlands. As with the impacts on tourism, impacts on all the latter sectors have to date only been assessed qualitatively.

4 Impacts on societal stability

The sea level is just one of several environmental factors that may be altered considerably by climate change. In combination with the other factors of climate change, it may have strong long-term impacts on the most downstream country of the Nile. A reduction in the amount of agricultural land available for food production and a decline in freshwater availability can force large numbers of people living in the coastal areas of Egypt to migrate to other parts of the countries. Such a development can increase the conflict potential within the Egyptian population and can thus contribute to a destabilization of the society as a whole.

But due to their different environmental characteristics, there needs to be a differentiation of the risks among the different coastal zones of Egypt. A qualitative assessment of the risks with regard to the consequences of sea level rise indicates that there are definite geographical differences (Tab. 2). The Nile Delta region faces the highest risks, and in all coastal sectors the economic risks from sea level rise are more prominent than the ecological risks or the direct risks for the population. Despite the relatively low effectiveness of coastal protection, it appears to be adequate for the current sea level (Frihy and El-Sayed,

2012). However, protective structures have to be maintained and even improved if the sea level rise accelerates. This requires substantial financial investments that may no longer be available if the country is affected by the combination of several climate change impacts such as increasing temperatures, even less precipitation in conjunction with a higher sea level and a possible increase in the frequency of storms.

Table 2: Risks with regard to sea level rise of the different sectors of the Egyptian Mediterranean coasts (adapted from Frihy and El-Sayed, 2012)

	population	economy	environment	effectiveness of protection
Northwest coast	low risk	moderate risk	low risk	low
Alexandria	moderate risk	high risk	moderate risk	moderate
Nile Delta	moderate risk	high risk	high risk	moderate
North Sinai	low risk	moderate risk	low risk	low




Figure 3: Schematic representation of the interactions between actors and issues in the Nile River Basin in times of climate change with a particular focus on the impacts of sea level rise (adapted from Scheffran et al., 2012c).

The impacts of climate change affect many parts of society and cause a multitude of responses. Some of them lead to increased cooperation and some may cause conflicts, possibly even violent conflicts (Link et al., 2012). These responses do not follow a simple pattern that can directly relate the occurrence of conflict to a particular change in environmental conditions. Instead, these responses are the result of the influence of the environment on the individuals living in the coastal zones of Egypt who then act and also interact with each other. These interactions between the various actors in the Nile River Basin and in this case particularly in Egypt are the basis of a model that has been developed at the Research Group Climate Change and Security (CLISEC) at the University of Hamburg (Scheffran et al., 2012b). It looks at the consequences of (climate change induced) environmental change for societal stability and considers the complex interactions between

climate change, environmental impacts, resource availability, individual human wellbeing and societal stability (Fig. 3).

Sea level rise is one of the factors directly influenced by climate change that has to be considered in this context as it aggravates the already difficult situation to adequately supply the Egyptian population with fundamental resources such as drinking water and fundamental crops. In the model framework, a combination of sea level rise, salinization of water resources, and land erosion affects the coastal zones of Egypt. This affects the land availability, mainly in the far downstream part of the Nile in the Nile Delta but also to a lesser extent in the eastern part of Egypt along the Red Sea coast. Land availability has an influence on agricultural and industrial production and also affects value added from tourism. Human wellbeing declines because of reduced economic revenues, which in turn further reduces the number of jobs available. Consequently, there is a self-reinforcing feedback loop back to economic output. Reduced job availability triggers out-migration into the other (presumably less affected) parts of the country, possibly upsetting the stability of the social systems and causing unrest and conflict.

More detailed assessments of the particular implications of sea level rise on the coastal zones of Egypt and their societal consequences are subject of future research to support this qualitative framework with results of quantitative model simulations. But the already existing results of initial assessments of the impacts of sea level rise on the Egyptian coastal zones indicate that large scale coastal protection is mandatory if the worst impacts, particularly due to inundation of the sea, are to be avoided.

5 Strategies to deal with sea level rise

There are already a number of measures taken to protect the Egyptian coast against the consequences of sea level rise. Already, more than US\$ 300 million have been spent to install sea walls along the most severely threatened parts of the coast (Elsharkawy et al., 2009). In some parts of the coast it is possible to rely on natural features with an elevation that is high enough to serve as protection (Frihy and El-Sayed, 2012). These include extensive sand dunes and ridges that run parallel to the shoreline. They are complemented by artificial structures such as seawalls or breakwaters. In particular, the outer parts of the promontories where the Nile River meets the sea are protected by seawalls. In these areas, erosion has declined considerably near the seawalls or has in some instances even changed to sand accumulation (Frihy et al., 2003). However, in the areas adjacent to these protective structures erosion has intensified in the downstream direction of the seawalls.

Prior to the Arab Spring in 2011 the Egyptian government had also started protecting the areas at risk by establishing new environmental regulations (Elsharkawy et al., 2009). First efforts to reduce the adverse impacts of sea level rise on water resources by implementing a National Improvement Plan are supposed to reach their targets by 2017. However, these plans have been criticized for mainly protecting tourism infrastructure. A more comprehensive protection plan would need to include stronger monitoring efforts and better enforcement of the measures. This includes the establishment of an Integrated Coastal Zone Management Committee and the improvement of community awareness of the risks of climate change to the coastal zones.

6 Conclusion

The current political situation in Egypt could prove to be a real problem when it comes to coastal protection. Assessments of the adaptive measures that have already been implemented indicate that these are currently insufficient to adequately prepare the Egyptian coastal zones for a future increase in sea level rise (Dasgupta et al., 2009a). But in times of

political turmoil and power changing hands, environmental protection often takes a back seat in comparison to individual rights and economic considerations of many. It is not clear how the importance of coastal protection will rank among the priorities of the many problems that need to be addressed by the new Egyptian administration in the coming years. Furthermore, it is traditionally not easy to implement plans for coastal protection as they require joint actions by all administrative levels in the country and a substantial increase in awareness of the expected problems that will arise in conjunction with sea level rise.

So despite the efforts to protect the Egyptian coasts against the impacts of sea level rise, particularly the coast along the Nile Delta is only protected in a very fragmented manner (Malm, 2012). This is mainly due to the limited resources available to Egypt to invest into adaptation measures to climate change and the fact that some tourism developments directly on the edge of the sea need particular protection. The protective measures for this infrastructure consume the majority of the available resources as considerable economic assets would be lost if these tourism resorts were abandoned. However, it is unlikely that coastal protection in Egypt can be successful if the prioritization in protection projects remains unaltered. If Egypt aims at successfully adapting to the consequences of sea level rise, it needs to abandon the business-as-usual protection strategies of the coasts and to implement a “revolutionary sea wall” (Malm, 2012:23) that includes the raising of public awareness and public participation which includes the whole society.

References

- Becker, R.H. and Sultan, M. 2009: Land subsidence in the Nile Delta: inferences from radar interferometry. *The Holocene* 19, 949-954.
- Church, J.A., Aarup, T., Woodworth, P.L., Wilson, W.S., Nicholls, R.J., Rayner, R., Lambeck, K., Mitchum, G.T., Steffen, K., Cazenave, A., Blewitt, G., Mitrovica, J.X. and Lowe, J.A. 2010: Sea-level Rise and Variability - Synthesis and Outlook for the Future. *Understanding Sea Level Rise and Variability*, Chichester, UK: Wiley-Blackwell, 402-420.
- Dasgupta, S., Laplante, B., Meisner, C., Wheeler, D. and Yan, J. 2009a: The impact of sea level rise on developing countries: a comparative analysis. *Climatic Change* 93, 379-388.
- Dasgupta, S., Laplante, B., Murray, S. and Wheeler, D. 2009b: Sea-Level Rise and Storm Surges: A Comparative Analysis of Impacts in Developing Countries. Washington, DC: The World Bank, 1-41.
- Dasgupta, S., Laplante, B., Murray, S. and Wheeler, D. 2011: Exposure of developing countries to sea-level rise and storm surges. *Climatic Change* 106, 567-579.
- El-Asmar, H.M. and Hereher, M.E. 2011: Change detection of the coastal zone east of the Nile Delta using remote sensing. *Environmental Earth Sciences* 62, 769-777.
- El Banna, M.M. and Frihy, O.E. 2009: Human-induced changes in the geomorphology of the northeastern coast of the Nile delta, Egypt. *Geomorphology* 107, 72-78.
- El Raey, M. 1997: Vulnerability assessment of the coastal zone of the Nile delta of Egypt, to the impacts of sea level rise. *Ocean & Coastal Management* 37, 29-40.
- El Raey, M. 2011: Mapping Areas Affected by Sea-Level Rise due to Climate Change in the Nile Delta Until 2100. In Brauch, H.G., Oswald-Spring, é., Mesjasz, C., Grin, J., Kameri-Mbote, P., Chourou, B., Dunay, P. and Birkmann, J., editors, *Coping with Global Environmental Change, Disasters and Security*, Heidelberg, Germany: Springer Verlag, 773-788.
- Elsharkawy, H., Rashed, H. and Rached, I. 2009: Climate Change: The Impacts of Sea Level Rise on Egypt. 1-11.
- FitzGerald, D.M., Fenster, M.S., Argow, B.A. and Buynevich, I.V. 2008: Coastal Impacts Due to Sea-Level Rise. *Annual Review of Earth and Planetary Sciences* 36, 601-647.

- Frihy, O.E. 2003: The Nile Delta-Alexandria Coast: Vulnerability to Sea-Level Rise, Consequences and Adaptation. *Mitigation and Adaptation Strategies for Global Change* 8, 115-138.
- Frihy, O.E., Deabes, E.A., Shereet, S.M. and Abdalla, F.A. 2010: Alexandria-Nile Delta coast, Egypt: update and future projection of sea-level rise. *Environmental Earth Sciences* 61, 253-273.
- Frihy, O.E., Debes, E.A. and El Sayed, W.R. 2003: Processes reshaping the Nile delta promontories of Egypt: pre- and post-protection. *Geomorphology* 53, 263-279.
- Frihy, O.E. and El-Sayed, M.K. 2012: Vulnerability risk assessment and adaptation to climate change induced sea level rise along the Mediterranean coast of Egypt. *Mitigation and Adaptation Strategies for Global Change* online first
- Frihy, O.E., Shereet, S.M. and El Banna, M.M. 2008: Pattern of Beach Erosion and Scour Depth along the Rosetta Promontory and their Effect on the Existing Protection Works, Nile Delta, Egypt. *Journal of Coastal Research* 24, 857-866.
- Hereher, M.E. 2010: Vulnerability of the Nile Delta to sea level rise: an assessment using remote sensing. *Geomatics, Natural Hazards and Risk* 1, 315-321.
- Intergovernmental Panel on Climate Change, editor 2007: *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge: Cambridge University Press.
- Jacob, T., Wahr, J., Pfeffer, W.T. and Swenson, S. 2012: Recent contributions of glaciers and ice caps to sea level rise. *Nature* 482, 514-518.
- Link, P.M., Piontek, F., Scheffran, J. and Schilling, J. 2012: On foes and flows: vulnerabilities, adaptive capacities and transboundary relations in the Nile River Basin in times of climate change. *Working paper CLISEC-19*, Hamburg, Germany: University of Hamburg, Research Group Climate Change and Security (CLISEC), 36.
- Malm, A. 2012: Sea Wall Politics: Uneven and Combined Protection of the Nile Delta Coastline in the Face of Sea Level Rise. *Critical Sociology*, 1 -30.
- McGranahan, G., Balk, D. and Anderson, B. 2007: The rising tide: assessing the risks of climate change and human settlements in low elevation coastal zones. *Environment & Urbanization* 19, 17-37.
- Nicholls, R.J., Hanson, S.E., Lowe, J.A., Warrick, R.A., Lu, X., Long, A.J. and Carter, T.R. 2011: Constructing Sea-Level Scenarios for Impact and Adaptation Assessment of Coastal Area: A Guidance Document. Intergovernmental Panel on Climate Change Task Group on Data and Scenario Support for Impacts and Climate Analysis (TGICA), 1-47.
- Rahmstorf, S. 2012: Modeling Sea Level Rise. *Nature Education Knowledge* 3, 4.
- Rahmstorf, S., Cazenave, A., Church, J.A., Hansen, J.E., Keeling, R.F., Parker, D.E. and Somerville, R.C.J. 2007: Recent Climate Observations Compared to Projections. *Science* 316, 709.
- Schaeffer, M., Hare, W., Rahmstorf, S. and Vermeer, M. 2012: Long-term sea-level rise implied by 1.5 °C and 2 °C warming levels. *Nature Climate Change* 2, 867-870.
- Scheffran, J., Brzoska, M., Kominek, J., Link, P.M. and Schilling, J. 2012a: Climate Change and Violent Conflict. *Science* 336, 869-871.
- Scheffran, J., Link, P.M. and Schilling, J. 2012b: Theories and Models of the Climate-Security Interaction: Framework and Application to a Climate Hot Spot in North Africa In Scheffran, J., Brzoska, M., Brauch, H.G., Link, P.M. and Schilling, J., editors, *Climate Change, Human Security and Violent Conflict*, Berlin: Springer, 91-131.
- Scheffran, J., Link, P.M. and Schilling, J. 2012c: Theories and Models of the Climate-Security Interaction: Framework and Application to a Climate Hot Spot in North Africa In Scheffran, J., Brzoska, M., Brauch, H.G., Link, P.M. and Schilling, J., editors, *Climate Change, Human Security and Violent Conflict: Challenges for Societal Stability*, Berlin: Springer, 91-131.
- Shaker, A.A., Alnaggar, D., Saad, A.A. and Faisal, H. 2011: Absolute Sea Level Rise Estimation at Alexandria Using Tide Records and GPS Observations. Marrakech, Morocco: FIG Working Week 2011, 1-14.

- Sherif, M.M. and Singh, V.J. 1999: Effect of climate change on sea water intrusion in coastal aquifers. *Hydrological Processes* 13, 1277-1287.
- Stammer, D. 2008: Response of the global ocean to Greenland and Antarctic ice melting. *Journal of Geophysical Research* 113, C06022.
- Stanley, D.L. 1998: Explaining persistent conflict among resource users: The case of Honduran mariculture. *Society & Natural Resources* 11, 267-278.
- Syvitski, J.P.M., Kettner, A.J., Overeem, I., Hutton, E.W.H., Hannon, M.T., Brakenridge, G.R., Day, J., Vörösmarty, C.J., Saito, Y., Giosan, L. and Nicholls, R.J. 2009: Sinking deltas due to human activities. *Nature Geoscience* 2, 681-686.
- Werner, A.D. and Simmons, C.T. 2009: Impact of Sea-Level Rise on Sea Water Intrusion in Coastal Aquifers. *Ground Water* 47, 197-204.

Acknowledgements

Research for this paper was supported through the Cluster of Excellence 'CliSAP' (EXC177), University of Hamburg, which is funded through the German Science Foundation (DFG).

CLISEC Working Papers

Scheffran, J. (2009): *Climate change, Human Security and Societal Instability: Conflict or Cooperation?*, Paper presented at the Synthesis Conference of the Global Environmental Change and Security program, Oslo, Norway, 22-24 June, 2009. Working paper **CLISEC-1**, Research Group Climate Change and Security, University of Hamburg.

Kominek, J. (2009): *Analysing the path of "using cyclical patterns for interpreting the situation" – a different approach using path dependency theory*, submitted. Working paper **CLISEC-2**, Research Group Climate Change and Security, University of Hamburg.

Scheffran, J., Link, P.M. & Schilling, J. (2009): *Theories and Models of the Climate Security Link*, Paper presented at the Conference "Climate Change, Social Stress and Violent Conflict", Hamburg, Germany, 19-20 November, 2009. Working paper **CLISEC-3**, Research Group Climate Change and Security, University of Hamburg.

Kominek, J. (2009): *Each institutionalization elementary is a self-reinforcing process increasing path dependency*, submitted. Working paper **CLISEC-4**, Research Group Climate Change and Security, University of Hamburg.

Kominek, J. (2009): *A new action model – deducing an 'ideal type path dependent' for scenario simulation*, submitted. Working paper **CLISEC-5**, Research Group Climate Change and Security, University of Hamburg.

Link, P.M., Piontek, F., Scheffran, J. & Schilling, J. (2010): *Integrated Assessment of Climate Security Hot Spots in the Mediterranean Region: Potential Water Conflicts in the Nile River Basin*, Paper presented at the Conference "Climate Change and Security", Trondheim, Norway, 21-24 June, 2010. Working paper **CLISEC-6**, Research Group Climate Change and Security, University of Hamburg.

Kominek, J. (2010): Global climate policy reinforces local social path dependent structures: More conflict in the world?, in: Scheffran, J., Brzoska, M., Brauch, H.G., Link, P.M. & Schilling, J., *Climate Change, Human Security and Violent Conflict: Challenges for Societal Stability*, Berlin, Springer Verlag, Hexagon Series Vol. 8, pp. 133-147. Working paper **CLISEC-7**, Research Group Climate Change and Security, University of Hamburg.

Kominek, J. (2010): *The Pursuit of Rational Action leads to Herding Behavior – an Example of Reinforcing Dynamics Shaped by Organizational Structures and Behavioral Paths in Financial Markets*, submitted. Working paper **CLISEC-8**, Research Group Climate Change and Security, University of Hamburg.

Link, P.M., Piontek, F., Scheffran, J. & Schilling, J. (2010): *Impact of climate change on water conflict and cooperation in the Nile River Basin*. Working paper **CLISEC-9**, Research Group Climate Change and Security, University of Hamburg.

Link, P.M. & Schleupner, C. (2010): How do tourists perceive and accept changes in landscape characteristics on the Eiderstedt peninsula?, *Coastline Reports*, **17**, pp. 133-146. Working paper **CLISEC-10**, Research Group Climate Change and Security, University of Hamburg.

Alwardt, C. (2011): *Wasser als globale Herausforderung: Die Ressource Wasser*. Working paper **CLISEC-11**, Research Group Climate Change and Security, University of Hamburg.

Kominek, J. & Scheffran, J. (2011): Cascading Processes and Path Dependency in Social Networks, in: Soeffner, H.-G. (ed.), *Transnationale Vergesellschaftungen*, Wiesbaden, VS Verlag für Sozialwissenschaften. Working paper **CLISEC-12**, Research Group Climate Change and Security, University of Hamburg.

Schilling, J., Freier, K., Hertig, E. & Scheffran, J. (2011): Climate change, vulnerability and adaptation in North Africa with focus on Morocco, *Agriculture, Ecosystems and Environment*, **156**, pp. 12-26. Working paper **CLISEC-13**, Research Group Climate Change and Security, University of Hamburg.

Schilling, J. & Remling, E. (2011): *Local Adaptation and National Climate Change Policy in Kenya: Discrepancies, Options, and the Way Forward*, submitted. Working paper **CLISEC-14**, Research Group Climate Change and Security, University of Hamburg.

Schilling, J., Akuno, M., Scheffran, J. & Weinzierl, T. (2011): *On arms and adaptation: Climate change and pastoral conflict in Northern Kenya*, submitted. Working paper **CLISEC-15**, Research Group Climate Change and Security, University of Hamburg.

Scheffran, J., Marmer, E. & Sow, P. (2011): Migration as a contribution for resilience and innovation in climate adaptation: Social networks and co-development in Northwest Africa, *Applied Geography*, **33**, pp. 119-127. Working paper **CLISEC-16**, Research Group Climate Change and Security, University of Hamburg.

Marmer, E., Scheffran, J. & Sow, P. (2011): *From security threat to conflict prevention: Integrating migration into climate adaptation policy frameworks in Africa*, submitted. Working paper **CLISEC-17**, Research Group Climate Change and Security, University of Hamburg.

Scheffran, J., Brzoska, M., Kominek, J., Link, P.M. & Schilling, J. (2012): Past and future research on climate change and violent conflict, *Review of European Studies*, **4** (5), pp. 1-13. Working paper **CLISEC-18**, Research Group Climate Change and Security, University of Hamburg.

Link, P.M., Piontek, F., Scheffran, J. & Schilling, J. (2012): *On foes and flows: Vulnerabilities, adaptive capacities and transboundary relations in the Nile River Basin in times of climate change*, submitted to *L'Europe en formation*. Working paper **CLISEC-19**, Research Group Climate Change and Security, University of Hamburg.

Link, P.M., Brzoska, M., Maas, A., Neuneck, G. & Scheffran, J. (2012): *Report on the conference "Geoengineering the Climate: An Issue for Peace and Security Studies?"*. Working paper **CLISEC-20**, Research Group Climate Change and Security, University of Hamburg.

Yang, L., Zhang, C. & Ngaruiya, G.W. (2012): Water Risks and Urban Responses under a Changing Climate: A Case Study of Hong Kong, *Pacific Geographies*, **39**, pp. 9-15. Working paper **CLISEC-21**, Research Group Climate Change and Security, University of Hamburg.

Kominek, J. & Link, P.M. (2012): *Modeling the linkage between climate change and violent conflict*, submitted. Working paper **CLISEC-22**, Research Group Climate Change and Security, University of Hamburg.

Ngaruiya, G. & Scheffran, J. (2012): *Reducing climate adaptation deficits using revolving fund network schemes in rural areas: Case study of Loitokitok district, Kenya*, submitted. Working paper **CLISEC-23**, Research Group Climate Change and Security, University of Hamburg.

Gioli, G. (2012): *Field Trip Report: Gender and Environmental Migration in the Karakoram Region*. Working paper **CLISEC-24**, Research Group Climate Change and Security, University of Hamburg.

Link, P.M., Kominek, J. & Scheffran, J. (2012): *Impacts of sea level rise on the coastal zones of Egypt*, submitted. Working paper **CLISEC-25**, Research Group Climate Change and Security, University of Hamburg.